

Two-point functions for $SU(3)$ Polyakov Loops near T_c

Adrian Dumitru and Robert D. Pisarski

Department of Physics, Brookhaven National Laboratory, Upton, New York 11973-5000, USA
emails: dumitru@quark.phy.bnl.gov and pisarski@quark.phy.bnl.gov

(February 1, 2008)

We discuss the behavior of two point functions for Polyakov loops in a $SU(3)$ gauge theory about the critical temperature, T_c . From a $Z(3)$ model, in mean field theory we obtain a prediction for the ratio of masses at T_c , extracted from correlation functions for the imaginary and real parts of the Polyakov loop. This ratio is $m_i/m_r = 3$ if the potential only includes terms up to quartic order in the Polyakov loop; its value changes as pentic and hexatic interctions become important. The Polyakov Loop Model then predicts how m_i/m_r changes above T_c .

Consider the behavior of an $SU(3)$ gauge theory, without dynamical quarks, at nonzero temperature. The usual quantity measured by the Lattice is the pressure. While this is all one needs for thermodynamics in equilibrium, near equilibrium how correlation functions behave is also of importance.

In this note we discuss a basic, gauge-invariant correlation function, the two-point function of Polyakov loops. We work with a $Z(3)$ model [1,2]; near the critical temperature, this is equivalent to the Polyakov Loop Model [3–6].

At a nonzero temperature T , the thermal Wilson line is:

$$\mathbf{L}(\vec{x}) = \mathcal{P} \exp \left(ig \int_0^{1/T} A_0(\vec{x}, \tau) d\tau \right), \quad (1)$$

with A_0 in the fundamental representation. This transforms as an adjoint field under the local $SU(3)/Z(3)$ gauge symmetry, and as a field with charge one under the global $Z(3)$ symmetry. To obtain a gauge invariant operator, the simplest thing to do is to take the trace, forming the Polyakov loop,

$$\ell = \frac{1}{3} \text{tr}(\mathbf{L}). \quad (2)$$

This transforms under the global $Z(3)$ symmetry as a field with charge one,

$$\ell \rightarrow \exp \left(\frac{2\pi i}{3} \right) \ell \quad (3)$$

The expectation value of ℓ is only nonzero above T_c , which is the temperature for the deconfining phase transition.

For three (or more) colors, ℓ is a complex valued field. Implicitly, we assume that the expectation value of ℓ is real. In this vacuum, we can then ask for the correlation functions of the real and imaginary parts of the Polyakov loop,

$$\ell_r = \text{Re } \ell, \quad \ell_i = \text{Im } \ell, \quad (4)$$

which do not mix. As the theory is globally $Z(3)$ invariant, we can define equivalent masses in either of the other two degenerate vacua by an appropriate overall $Z(3)$ rotation.

We need a standard feature of propagators in coordinate space. In d space-time dimensions, at distances large compared to $1/T$, only the $d-1$ spatial dimensions matter. For a field of mass m , at distances $x \gg 1/m$, of course the propagator falls off exponentially,

$$\int d^{d-1}p \frac{e^{-i\vec{p}\cdot\vec{x}}}{p^2 + m^2} \sim \int d^{d-2}p_\perp e^{-x(m+p_\perp^2/2m)} \sim \frac{e^{-mx}}{x^{(d-2)/2}}. \quad (5)$$

We are interested not in the familiar exponential fall-off, but in power law prefactor of $1/x^{(d-2)/2}$. While derived in free field theory, it is valid generally. It arises from the fluctuations of a point particle, in the directions perpendicular to the direction along which we measure it.

In perturbation theory, the Polyakov loop is near unity, and can be expanded in powers of A_0 . Assuming for the purposes of discussion that the A_0 field is purely static, we obtain

$$\ell \approx 1 - \frac{g^2}{3T^2} \text{tr}(A_0^2) - i \frac{g^3}{3T^3} \text{tr}(A_0^3) + \dots \quad (6)$$

Thus the real part of the Polyakov loop starts out as a coupling to two A_0 's, and the imaginary part of the Polyakov loop, to three A_0 's. Thus to lowest order in perturbation theory [7],

$$\langle \ell_r(x) \ell_r(0) \rangle - \langle \ell \rangle^2 \sim \frac{\exp(-2m_D x)}{x^2}, \quad (7)$$

$$\langle \ell_i(x) \ell_i(0) \rangle \sim \frac{\exp(-3m_D x)}{x^3}. \quad (8)$$

Here m_D is the Debye mass; to lowest order in perturbation theory, $m_D = gT$. Each result is simply the product of either two or three A_0 propagators over distances $x \gg 1/m_D$. Thus, if one ignores the difference

in prefactors, and defines m_r and m_i from the exponential falloff of (7) and (8), respectively, we conclude that $m_i/m_r = 3/2$. These estimates could be altered by mixing with the purely magnetic sector [8], which we discuss at the end of the paper.

We now turn from the perturbative regime, valid at very high temperatures, to temperatures near T_c . Instead of dealing with A_0 , we construct an effective Lagrangian directly for the Polyakov loop itself [1,2]. The theory must be invariant under the global $Z(3)$ symmetry of (3). Implicitly, we assume that the deconfining transition for three colors is so weakly first order that it is appropriate to classify the effective Lagrangian according to a renormalization group analysis for a second order transition. At large distances, $x \gg 1/T$, the theory is effectively three dimensional; by the renormalization group, all relevant and marginal operators include terms up to sixth order in ℓ . The most general, renormalizable Lagrangian in three dimensions, invariant under a global $Z(3)$ symmetry, is

$$\begin{aligned} \mathcal{Z}_W |\vec{\partial}\ell|^2 - \frac{b_2}{2} |\ell|^2 - \frac{b_3}{3} \left(\frac{\ell^3 + (\ell^*)^3}{2} \right) \\ + \frac{b_4}{4} (|\ell|^2)^2 + \frac{b_5}{5} |\ell|^2 \left(\frac{\ell^3 + (\ell^*)^3}{2} \right) \\ + \frac{b_6}{6} (|\ell|^2)^3 + \frac{\tilde{b}_6}{6} \left(\frac{\ell^3 + (\ell^*)^3}{2} \right)^2. \end{aligned} \quad (9)$$

The terms which only involve powers of $|\ell|^2$, proportional to $\sim b_2$, b_4 and b_6 , are each each invariant under a global symmetry of $U(1)$, $\ell \rightarrow \exp(i\theta)\ell$ for arbitrary θ . The kinetic term is also $U(1)$ invariant. For reasons to become clear shortly, we include a factor of wave-function renormalization, $\mathcal{Z}_W$, but suppress all other renormalization constants for mass and coupling constants. The cubic term, $\sim b_3 \ell^3$, and the pentic term, $\sim b_5 |\ell|^2 \ell^3$, are only invariant under the $Z(3)$ symmetry of (3). Of the two hexatic interactions, that $\sim b_6$ is $U(1)$ invariant, while that $\sim \tilde{b}_6 \ell^6$ is invariant not only under $Z(3)$, but under a larger symmetry of $Z(6)$, $\ell \rightarrow \exp(\pi i/3)\ell$.

It is natural to require that the potential is bounded from below for large values of ℓ . At large ℓ , the hexatic terms dominate over all other couplings. For real ℓ , this gives

$$b_6 + \tilde{b}_6 > 0. \quad (10)$$

One can also choose $\ell = \exp(i\pi/6)\ell_0$, with real ℓ_0 ; then the term proportional to $\tilde{b}_6$ drops out, giving the condition

$$b_6 > 0. \quad (11)$$

We perform a mean field analysis, where all coupling constants are taken as constant with temperature, except for the mass term, $\sim b_2 |\ell|^2$. About the transition,

condensation of ℓ is driven by changing the sign of b_2 [2,3]. The sign of b_3 is determined by the following. At very high temperature, the favored vacuum is perturbative, with $\ell_0 \approx 1$, times $Z(3)$ rotations. We then choose $b_3 > 0$ so that in the $Z(3)$ model, there is always one vacuum with a real, positive expectation value for ℓ_0 .

We also choose $b_4 > 0$. This is not a matter of convention. With the above potential, in mean field theory there are two ways of obtaining a first order transition. One is through a non-zero, cubic coupling. The other is if the four point coupling is negative. It is natural, however, to assume that $b_3, b_5 \neq 0$ and $b_4 > 0$. For example, if $b_3 = b_5 = 0$, then the unbroken symmetry of the effective Lagrangian is not $Z(3)$, but $Z(6)$. While one can then obtain a first order transition by choosing b_4 to be negative, it would also imply a six-fold degeneracy between vacua in the broken symmetry phase. Such a degeneracy is not observed, even approximately, by Lattice simulations in the deconfined phase [9]. Consequently, we assume that $b_3, b_5 \neq 0$, and that b_4 is positive. This also agrees with results from the Polyakov Loop Model [3,4].

While the value of the pentic coupling constant is not well fixed, but it cannot dominate the hexatic couplings. If the hexatic interactions vanish, $b_6 = \tilde{b}_6 = 0$, then with $\ell = \exp(i\theta/3)\ell_0$, the pentic term is negative for $\pi/2 < \theta < 3\pi/2$, and the potential is unbounded from below. Consequently, we consider the pentic interactions as a small perturbation on the other couplings.

This is an effective Lagrangian in three dimensions, so that ℓ has dimensions of mass^{1/2}; then the coupling constants b_n have dimensions of mass^{(6-n)/2}. In particular, b_6 and $\tilde{b}_6$ are dimensionless. To make the equations somewhat simpler, we choose the overall mass scale so that $b_4 = 1$; expressions for arbitrary values of b_4 can be derived from the following by an obvious rescaling. We remark that in the Polyakov Loop Model [3–6], the dimensions of ℓ , and of the coupling constants, are all made up by powers of temperature. Near T_c , however, this doesn't matter.

We now compute the masses for the real and imaginary parts of the Polyakov loop. As stated, we choose the vacuum expectation value of ℓ , $\langle \ell \rangle = \ell_0$, to be real. The equation for ℓ_0 is just the first derivative of the potential for ℓ ,

$$-b_2 \ell_0 - b_3 \ell_0^2 + \ell_0^3 + b_5 \ell_0^4 + (b_6 + \tilde{b}_6) \ell_0^5 = 0. \quad (12)$$

Computing second derivatives of the potential, the mass squared for the real part is:

$$m_r^2 = -b_2 - 2b_3 \ell_0 + 3\ell_0^2 + 4b_5 \ell_0^3 + 5(b_6 + \tilde{b}_6) \ell_0^4, \quad (13)$$

while that for the imaginary part is:

$$m_i^2 = -b_2 + 2b_3 \ell_0 + \ell_0^2 - \frac{4}{5} b_5 \ell_0^3 + (b_6 - 2\tilde{b}_6) \ell_0^4. \quad (14)$$

The two-point function of Polyakov loops is then,

$$\langle \ell_r(x) \ell_r(0) \rangle - \langle \ell \rangle^2 \sim \frac{\exp(-m_r x)}{x}, \quad (15)$$

$$\langle \ell_i(x) \ell_i(0) \rangle \sim \frac{\exp(-m_i x)}{x}. \quad (16)$$

Notice that unlike (7) and (8), the power law prefactor is in each case just $\sim 1/x$, from the exchange of a single ℓ -field. Above T_c , $m_i \neq m_r$.

The global $Z(3)$ symmetry is unbroken below T_c : $\ell_0 = 0$. With our conventions, $b_2 < 0$; the mass squared for ℓ_r and ℓ_i are positive and equal. Thus in our $Z(3)$ model, below T_c the two-point functions of Polyakov loops remain as in (15) and (16) at large distances, with $m_i = m_r$. We remark that the prefactor of $\sim 1/x$, for $T < T_c$ and $x \gg 1/T$, agrees with calculations in Nambu-Goto string models [10]. This is true for any number of space-time dimensions: the transverse motion of a point particle, and the transverse fluctuations of a world-sheet, both give the same prefactor of $\sim 1/x^{(d-2)/2}$. This is an important consistency check on models of Polyakov loops, such as [3], below T_c .

When b_3 or b_5 are nonzero, the transition is necessarily of first order. Henceforth, we restrict ourselves to the critical temperature, T_c . This occurs when the nontrivial minimum is degenerate with the trivial minimum; *i.e.*, when

$$-\frac{b_2}{2} \ell_0^2 - \frac{b_3}{3} \ell_0^3 + \frac{1}{4} \ell_0^4 + \frac{b_5}{5} \ell_0^5 + \frac{b_6 + \tilde{b}_6}{6} \ell_0^6 = 0. \quad (17)$$

This then becomes an equation which determines the value of $b_2(T_c^+)$.

If we compute each mass individually, m_r and m_i , then there is a complication which we have set aside. The kinetic term for ℓ does not have a standard normalization; at tree level, it equals $3/g^2$, where g is the QCD coupling constant [3]. At one loop order, Wirstam [5] finds that this wave function renormalization is

$$\mathcal{Z}_W = \frac{3}{g(T)^2} \left(1 - .08 \frac{g^2}{4\pi} + \dots \right). \quad (18)$$

Consequently, the masses measured on the Lattice are actually $1/\sqrt{\mathcal{Z}_W}$ of the expressions above. Thus the overall scale of m_r and m_i depend upon the value of $g^2(T)$, which is not easy to extract directly from Lattice data.

To avoid the dependence on $\mathcal{Z}_W$, instead we compute the ratio of m_i/m_r , in which $\mathcal{Z}_W$ completely drops out. It is this ratio which we think would be especially interesting to compute on the Lattice.

These equations can be solved numerically for arbitrary values of b_5 , b_6 , and $\tilde{b}_6$. It is illustrative to first solve for the case in which $b_5 = b_6 = \tilde{b}_6 = 0$. For arbitrary b_2 ,

$$\ell_0 = \frac{1}{2} \left(b_3 + \sqrt{4b_2 + b_3^2} \right). \quad (19)$$

With this ℓ_0 , at the transition temperature, $b_2 = -2b_3^2/9$, and $m_i/m_r = 3$. This ratio changes as b_5 , b_6 , and $\tilde{b}_6$ increase from zero. To lowest nontrivial order in these coupling constants,

$$\frac{m_i}{m_r} \approx 3 \left(1 - \frac{8}{15} b_3 b_5 - \frac{4}{27} b_3^2 (4b_6 + 5\tilde{b}_6) \right) + \dots \quad (20)$$

For positive values of b_5 and $\tilde{b}_6$ (remember b_6 is necessarily positive), the ratio m_i/m_r decreases from 3 as b_5 , b_6 , and $\tilde{b}_6$ increase. It is also easy to obtain $m_i/m_r > 3$, though. The coupling b_5 could be negative; even if $b_5 = 0$, when $\tilde{b}_6 = -b_6$, *etc.*

One can also solve for large values of b_6 and $\tilde{b}_6$. Taking $b_5 = 0$, in this case

$$\frac{m_i}{m_r} \approx \sqrt{\frac{3}{2}} \sqrt{1 + \frac{b_6}{b_6 + \tilde{b}_6}} + \dots \quad (21)$$

The combination of b_6 and $\tilde{b}_6$ on the right hand side is necessarily a ratio of positive numbers in all cases.

Working up from T_c , it is also possible to compute m_i/m_r within the context of the Polyakov Loop Model (PLM) [3]. In the PLM, the pressure is determined by the potential of ℓ times T^4 ; for constant values of b_3 , b_4 , b_5 , b_6 , and $\tilde{b}_6$, the temperature dependence of the pressure then fixes $b_2(T)$. Consequently, this also determines how m_i/m_r changes with temperature. This appears to be a crucial and unambiguous test of the PLM. For example, if we take $b_6 = \tilde{b}_6 = 0$, then at very high temperatures, where $\ell_0 \approx 1$, (19) fixes $b_2 = 1 - b_3$, and [6]

$$\frac{m_i}{m_r} = \sqrt{\frac{3b_3}{2 - b_3}}. \quad (22)$$

To agree with the perturbative result as $T \rightarrow \infty$, $= 3/2$, fixes $b_3 = 6/7$. This is remarkably close to the value found from a fit to the pressure near T_c , which is $b_3 \approx 0.9$ [11]. A more careful analysis, including the effects of pentic and hexatic interactions, is necessary in order to make a detailed comparison.

To summarize, naively in the perturbative regime at very high temperature, $m_i/m_r = 3/2$ [7]. If pentic and hexatic terms in ℓ are a small perturbation on the quartic terms, then as $T \rightarrow T_c$, m_i/m_r should increase; at T_c , how close this ratio is to 3 then indicates how important these other coupling constants are. Conversely, if m_i/m_r does not increase significantly as the temperature decreases from $2T_c$ down to T_c^+ , then while it might be able to obtain a fit with large values of b_6 and $\tilde{b}_6$, to us this would seem unnatural, and disfavor the present analysis.

Our discussion neglects mixing between the electric sector and the static, magnetic sector [8]. While the former might be perturbative, the latter never is. Static, magnetic glueballs couple to each other as a three-dimensional gauge theory, with coupling constant g^2T . Hence the mass of any static, magnetic glueball is some number times g^2T . In weak coupling, then, as the Debye mass is $m_D \sim gT$, electric fields are much heavier than magnetic glueballs. The real part of the Polyakov loop, $\sim \text{tr}(A_0^2)$, can then mix with spin-zero glueballs, such as $\sim \text{tr}(G_{ij}^2)$. Also, the imaginary part of the Polyakov loop, $\sim \text{tr}(A_0^3)$ can mix with the spin-zero glueball $\text{tr}(A_0 G_{ij}^2)$. (There is also a spin-one glueball, $\epsilon^{ijk} \text{tr}(A_0 G_{jk})$. However, the coupling of such a spin-one state to any spin-zero state must involve one derivative, and so vanish at zero momentum. Spin here refers to a classification in three dimensions.)

Simulations from temperatures of $\sim 2T_c$ on up have been done by Hart, Laine, and Philipsen [12]; for related work, see [13]. They find that the magnetic sector is not lighter than the electric until temperatures $T > 10^2 T_c$. Unexpectedly, they do not see significant mixing between states such as $\text{tr}(A_0^2)$ and $\text{tr}(G_{ij}^2)$. For example, at extremely high temperatures of $10^{11} T_c$, the mass of $\text{tr}(A_0^2)$ is approximately equal to that of $\text{tr}(A_0^3)$; both masses are about ≈ 1.9 times the mass of the lightest static, magnetic glueball, $\text{tr}(G_{ij}^2)$. The lightest electric state is the spin-one glueball, $\epsilon^{ijk} \text{tr}(A_0 G_{jk})$; we note the (presumably meaningless) numerical coincidence that its mass is $= 1.5$ of the lightest static, magnetic glueball to within 0.1%, although the quoted statistical errors are $\approx \pm 6\%$.

At temperatures of $\sim 2T_c$, the situation is completely reversed: then the electric sector is *lighter* than that for the static, magnetic sector. The lightest state is that for the real part of the Polyakov loop, $\sim \text{tr}(A_0^2)$. Relative to this state, the mass of the spin-zero electric state, $\text{tr}(A_0^3)$, is ≈ 1.81 ; that of the spin-one electric state, $\epsilon^{ijk} \text{tr}(A_0 G_{jk})$, is ≈ 1.76 ; that of the lightest static, magnetic glueball, $\text{tr}(G_{ij}^2)$, is ≈ 2.4 .

Thus at $2T_c$, the ratio $m_i/m_r \approx 1.81$. It would be very interesting to know how m_i/m_r changes at $T : 2T_c \rightarrow T_c$; we are unaware of data on m_i in this range. Nevertheless, at least we can safely neglect mixing of the (light) electric sector with the (heavy) magnetic sector.

At present, most Lattice data is for the two point function of complex conjugate Polyakov loops, $\langle \ell(x) \ell^*(0) \rangle - |\langle \ell \rangle|^2$. From either the $Z(3)$ model, or perturbation theory, in the deconfined phase it appears that m_i is always greater than m_r . Thus without extracting excited states, this correlation function only gives us information on m_r , and not on m_i .

One can also decide whether an effective model of Polyakov loops applies by seeing if the power law prefactor is $1/x$, as in (15), or $1/x^2$, as in (7). From Lattice simulations in $3+1$ dimensions by Kaczmarek, Karsch, Laermann, and Lutgemeier [14], the power law prefactor is very near unity only near T_c , and increases to ~ 1.4

by $2T_c$. Thus our effective model does appear to hold in at least some limited region near T_c . As the exchange of *any* massive state generates a prefactor of $\sim 1/x$, instead of varying the prefactor, to us it seems more natural to take a sum over several states, each a prefactor of $\sim 1/x$. We do admit that this is only true in mean field theory, as (nearly) critical fluctuations introduce nonzero anomalous dimensions, η , which change the prefactor to $\sim 1/x^{1-\eta}$. In three dimensions, however, such anomalous dimensions η are usually very small; $\eta < .1$ for either $Z(2)$ or $O(2)$ symmetry groups [15].

It is also of interest to consider the transition in $2+1$ dimensions. While mean field theory suggests a first order transition, fluctuations in two dimensions are so strong that the transition is in fact of second order. Then our mean field analysis is completely inappropriate; one is at the symmetric phase at T_c , with $m_i = m_r$. However, correlation functions of both imaginary and real parts of the Polyakov loop have been computed above T_c by Bialas, Morel, Petersson, and Petrov [16]. In all cases, the power law prefactor is found to be $1/\sqrt{x}$, which is what it should be for single particle exchange in $d = 2 + 1$ dimensions. Bialas *et al.* have also computed m_i/m_r ; working down from high temperatures, this ratio increases to ≈ 2.0 , then seems to decrease. This might arise from a trade-off between mean field effects, as discussed herein, and critical fluctuations near T_c , special to $2 + 1$ dimensions.

In passing, we also comment that studies with two colors are also of interest. For two colors, ℓ is real, and so there is only m_r . One could test, however, to see whether the relation between m_r and the pressure, as given by the PLM, holds up [3]. With only one mass to measure, the value of m_r does depend upon Z_W , and so $g^2(T)$. Also, the effects of critical fluctuations must be incorporated. On the other hand, the potential for ℓ only involves two coupling constants, b_4 , and b_6 ; $b_3 = b_5 = \tilde{b}_6 = 0$. Still, one basic test is that in $3 + 1$ dimensions, the deduced value of $g^2(T)$ increases as T decreases. Lastly, there are speculative reasons for thinking that the width of the critical region is greater in $2 + 1$ dimensions than in $3 + 1$ dimensions [2].

We look forward to future measurements of these and other quantities on the Lattice, near the critical temperature.

Acknowledgements: This research was supported by DOE grant DE-AC02-98CH10886. We thank A. D. Hart, J. T. Lenaghan, S. Ohta, and O. Philipsen for useful comments and discussions. We are especially indebted to M. Laine for pointing out the existence of the pentic coupling.

[1] B. Svetitsky and L. G. Yaffe, Nucl. Phys. B **210**, 423 (1982).

- [2] R. D. Pisarski, arXiv:hep-ph/0203271.
- [3] R. D. Pisarski, Phys. Rev. D **62**, 111501 (2000); A. Dumitru and R. D. Pisarski, Phys. Lett. B **504**, 282 (2001); A. Dumitru, O. Scavenius, and A. D. Jackson, Phys. Rev. Lett. **87**, 182302 (2001); O. Scavenius, A. Dumitru and J. T. Lenaghan, arXiv:hep-ph/0201079; P. N. Meisinger, T. R. Miller, and M. C. Ogilvie, arXiv:hep-ph/0108009; P. N. Meisinger and M. C. Ogilvie, arXiv:hep-ph/0108026.
- [4] A. Dumitru and R. D. Pisarski, Phys. Lett. B **525**, 95 (2002).
- [5] J. Wirstam, Phys. Rev. D **65**, 014020 (2002).
- [6] A. Dumitru and R. D. Pisarski, Nucl. Phys. Proc. Suppl. B **106**, 483 (2002).
- [7] S. Nadkarni, Phys. Rev. D **33**, 3738 (1986); *ibid.* **34**, 3904 (1986).
- [8] A. K. Rebhan, Phys. Rev. D **48**, 3967 (1993); Nucl. Phys. B **430**, 319 (1994); E. Braaten and A. Nieto, Phys. Rev. Lett. **74**, 3530 (1995); Phys. Rev. D **51**, 6990 (1995); P. Arnold and L. G. Yaffe, *ibid.* **52**, 7208 (1995).
- [9] T. A. DeGrand and C. E. DeTar, Phys. Rev. D **34**, 2469 (1986); P. Bacilieri *et al.*, Phys. Rev. Lett. **61**, 1545 (1988); F. R. Brown *et al.*, *ibid.* **61**, 2058 (1988); A. Ukawa, Nucl. Phys. Proc. Suppl. B **17**, 118 (1990); *ibid.* **53**, 106 (1997); E. Laermann, *ibid.* **63**, 114 (1998); F. Karsch, *ibid.* **83**, 14 (2000).
- [10] P. de Forcrand, G. Schierholz, H. Schneider and M. Teper, Phys. Lett. B **160**, 137 (1985).
- [11] This value of b_3 follows from Scavenius, Dumitru, and Lenaghan, [3]. This value follows not only by fitting the potential for ℓ to the pressure, but also requiring that $\ell_0 \rightarrow 1$ as $T \rightarrow \infty$.
- [12] A. Hart, M. Laine and O. Philipsen, Nucl. Phys. B **586**, 443 (2000).
- [13] M. Laine and O. Philipsen, Nucl. Phys. B **523**, 267 (1998); Phys. Lett. B **459**, 259 (1999); S. Datta and S. Gupta, Nucl. Phys. B **534**, 392 (1998); Phys. Lett. B **471**, 382 (2000); A. Hart and O. Philipsen, Nucl. Phys. B **572**, 243 (2000); R. V. Gavai and S. Gupta, Phys. Rev. Lett. **85**, 2068 (2000).
- [14] O. Kaczmarek, F. Karsch, E. Laermann and M. Lutgemeier, Phys. Rev. D **62**, 034021 (2000).
- [15] J. Zinn-Justin, "Quantum Field Theory and Critical Phenomena". (Oxford, Oxford University Press, 1989).
- [16] P. Bialas, A. Morel, B. Petersson and K. Petrov, Nucl. Phys. Proc. Suppl. B **106**, 882 (2002).